

Speed of convergence of two-dimensional Fourier integrals

Mark A Pinsky , Northwestern University

1. Introduction

Recently [2,3] we found necessary and sufficient conditions for the convergence at a preassigned point of the spherical partial sums of the Fourier integral in a class of piecewise smooth functions in Euclidean space. These yield elementary examples of divergent Fourier integrals in three dimensions and higher. Meanwhile, several years ago Gottlieb and Orsag[1] observed that in two dimensions we may expect slower convergence at certain points, specifically for Fourier-Bessel series of radial functions.

In this paper we investigate the rate of convergence of the spherical partial sums of the Fourier integral for a class of piecewise smooth functions. The basic result is an asymptotic expansion which allows us to read off the rate of convergence at a pre-assigned point.

2. Statement of results

$\mathbf{R}^2$ denotes the Euclidean plane and $f \in L^1(\mathbf{R}^2)$ is an integrable function with Fourier transform and spherical partial sum denoted by

$$\hat{f}(\xi_1, \xi_2) = \frac{1}{(2\pi)^2} \int_{\mathbf{R}^2} f(x_1, x_2) e^{-i(\xi_1 x_1 + \xi_2 x_2)} dx_1 dx_2,$$
$$f_M(x_1, x_2) = \int_{\xi_1^2 + \xi_2^2 \leq M^2} \hat{f}(\xi_1, \xi_2) e^{i(\xi_1 x_1 + \xi_2 x_2)} d\xi_1 d\xi_2.$$

The spherical mean value with respect to $x = (x_1, x_2)$ is defined by

$$\bar{f}_x(r) = \frac{1}{2\pi} \int_0^{2\pi} f(x_1 + r \cos \theta, x_2 + r \sin \theta) d\theta.$$

We say that f is piecewise smooth of class C^k with respect to $x \in \mathbf{R}^2$ iff the mapping $r \rightarrow \bar{f}_x(r)$ is piecewise

C^k and vanishes for $r \geq r(x)$. Such a function has left and right limits at all points, together with the requisite derivatives. These will agree except at a finite set of points which we denote by $a_i, 1 \leq i \leq K$. The jumps are denoted by

$$\delta \bar{f}_x(a_i) := \bar{f}_x(a_i + 0) - \bar{f}_x(a_i - 0),$$

with a corresponding notation for the jumps of the derivatives.

Theorem Suppose that f is piecewise smooth of class C^2 with respect to $x \in \mathbf{R}^2$. Then we have the asymptotic ($M \rightarrow \infty$) formula

$$f_M(x) - \bar{f}_x(0+0) = \sqrt{\frac{2}{M\pi}} \sum_{1 \leq i \leq K} \delta \bar{f}_x(a_i) \frac{\cos(Ma_i - \pi/4)}{\sqrt{a_i}} \\ + \frac{1}{M} \bar{f}'_x(0+0) + o(1/M).$$

Corollary *If the spherical mean value with respect to x is continuous, then we have*

$$f_M(x) - \bar{f}_x(0+0) = O(1/M), \quad M \rightarrow \infty.$$

Otherwise

$$-\infty < \liminf_M \sqrt{M}[f_M(x) - \bar{f}_x(0+0)] < \limsup_M \sqrt{M}[f_M(x) - \bar{f}_x(0+0)] < \infty.$$

Example If we choose $f(x) = 1_{[0,R]}(|x|)$ as the indicator function of a disc of radius R centered at $(0,0)$, then $\delta \bar{f}_x(r) = 0$ unless $x = 0, r = R$, where $\delta \bar{f}_0(R) = -1$; thus

$$f_M(0,0) = 1 - \sqrt{\frac{2}{M\pi R}} \cos(MR - \pi/4) + O(1/M), \\ f_M(x_1, x_2) = 1 + O(1/M) \quad 0 < x_1^2 + x_2^2 < R^2, \\ f_M(x_1, x_2) = \frac{1}{2} + O(1/M) \quad x_1^2 + x_2^2 = R^2, \\ f_M(x_1, x_2) = O(1/M) \quad x_1^2 + x_2^2 > R^2.$$

The speed of convergence at the center is strictly slower than at all other points.

3. Proofs

The spherical partial sum is written directly in terms of the spherical mean value by writing [3]

$$f_M(x) = \int_0^a M J_1(Mr) \bar{f}_x(r) dr, \quad a = r(x).$$

From the identities for Bessel functions, we have

$$M J_1(Mr) = -\frac{d}{dr} J_0(Mr),$$

so that we may integrate by parts:

$$f_M(x) = - \int_0^a \bar{f}_x(r) \frac{d}{dr} J_0(Mr) dr \\ = \bar{f}_x(0+0) + \sum_{1 \leq i \leq K} \delta \bar{f}_x(a_i) J_0(Ma_i) + \int_0^a J_0(Mr) \bar{f}'_x(r) dr.$$

The Bessel function has the asymptotic behavior $J_0(x) = \sqrt{(2/\pi x)}[\cos(x - \pi/4) + O(1/x)]$, $x \rightarrow \infty$, which yields the first term. To handle the integral term, we define $K(x) = \int_0^x J_0(t) dt$. From the asymptotic behavior of J_0 it follows that $K(x)$ is bounded with a limit $K(\infty) = \int_0^\infty J_0(t) dt = 1$ (see the proof in the appendix below). Therefore we may integrate by parts once again to obtain

$$\begin{aligned} \int_0^a J_0(Mr) \bar{f}'_x(r) dr &= \frac{1}{M} \int_0^a \bar{f}'_x(r) \frac{d}{dr} K(Mr) dr \\ &= -\frac{1}{M} \left(\sum_{1 \leq i \leq K} \delta \bar{f}'_x(a_i) K(Ma_i) + \int_0^a \bar{f}''_x(r) K(Mr) dr \right). \end{aligned}$$

When $M \rightarrow \infty$ the first set of terms within the parenthesis clearly converges to $K(\infty) \sum_{1 \leq i \leq K} \delta \bar{f}'_x(a_i)$; the second term converges, by the dominated convergence theorem, to $K(\infty) \int_0^a \bar{f}''_x(r) dr$. Applying the fundamental theorem of calculus on each sub-interval (a_{i-1}, a_i) and collecting terms gives the stated form. The proof is complete.

4. Comparison with one-dimensional convergence

In order to put the above in some perspective, we outline here a treatment of the corresponding questions for Fourier integrals in one variable, where we find a generic rate of convergence of $1/M$, even in the absence of continuity.

An integrable function $f(x)$, $x \in \mathbf{R}^1$ has the Fourier transform $\hat{f}(\xi) = (2\pi)^{-1} \int_{\mathbf{R}^1} f(x) e^{-i\xi x} dx$ with spherical partial sum $f_M(x) = \int_{-M}^M \hat{f}(\xi) e^{i\xi x} d\xi$. This can be expressed in terms of the *symmetric average*

$$\bar{f}_x(t) = \frac{1}{2} [f(x+t) + f(x-t)] \text{ as}$$

$$f_M(x) = \frac{2}{\pi} \int_0^\infty \frac{\sin Mt}{t} \bar{f}_x(t) dt$$

Using this representation, we now state and prove

Proposition. Suppose that $t \rightarrow \bar{f}_x(t)$ is piecewise C^2 and zero for $t > a$. Then we have the asymptotic formula

$$f_M(x) - \bar{f}_x(0+0) = \frac{2}{M\pi} [\bar{f}'_x(0+0) + \sum_{1 \leq i \leq K} \delta \bar{f}_x(a_i) \frac{\cos Ma_i}{a_i}] + o(1/M).$$

Proof. We write

$$f_M(x) - \bar{f}_x(0+0) = \frac{2}{\pi} \int_0^a \frac{\bar{f}_x(t) - \bar{f}_x(0+0)}{t} \sin Mt dt - \frac{2}{\pi} \bar{f}_x(0+0) \int_a^\infty \frac{\sin Mt}{t} dt,$$

where the last integral is conditionally convergent. In the first integral we write $g(t) = \frac{\bar{f}_x(t) - \bar{f}_x(0+0)}{t}$ and integrate by parts in the form

$$\begin{aligned} \int_0^a g(t) \sin Mt \, dt &= -(1/M) \int_0^a g(t) d[\cos Mt] \\ &= (1/M)[g(0) + \sum_{1 \leq i \leq K} \delta g(a_i) \cos Ma_i + \int_0^a g'(t) \cos Mt \, dt]. \end{aligned}$$

But $\delta g(a_i) = \frac{\delta \bar{f}_x(a_i)}{a_i}$.

The second integral can also be integrated by parts to obtain the asymptotic form

$$\int_a^\infty \frac{\sin Mt}{t} \, dt = \frac{\cos Ma}{Ma} + O(1/M^2).$$

Combining these produces the stated result.

Remark. A parallel result hold for Fourier series on the circle $-\pi < x < \pi$. In this case the spherical partial sum of $2M = 2n + 1$ terms is written

$$f_M(x) = \frac{2}{\pi} \int_0^\pi \frac{\sin Mt}{2 \sin(t/2)} \bar{f}_x(t) \, dt.$$

The above computations can be replicated with the result

$$f_M(x) - \bar{f}_x(0+0) = \frac{2}{M\pi} [\bar{f}'_x(0+0) + \sum_{1 \leq i \leq K} \delta \bar{f}_x(a_i) \frac{\cos Ma_i}{2 \sin(a_i/2)}] + o(1/M)$$

Conclusion. We note that the one-dimensional result (for both Fourier series and integrals) differs from the two-dimensional result in the appearance of the derivative term $\bar{f}'_x(0+0)$ at the *very first level* of the asymptotic analysis.

5. Appendix. Computation of $K(\infty)$

The Bessel function can be written

$$J_0(x) = \frac{1}{\pi} \int_{-1}^1 \frac{\cos xt}{\sqrt{(1-t^2)}} \, dt.$$

Thus by the Fubini theorem

$$\int_0^M J_0(x) \, dx = \frac{1}{\pi} \int_{-1}^1 \frac{\sin Mt}{t} \frac{1}{\sqrt{(1-t^2)}} \, dt.$$

This is the partial Fourier integral of the integrable function $(1-t^2)^{-1/2}$ at $t = 0$.

When $M \rightarrow \infty$ we can appeal to the convergence of one-dimensional Fourier integrals to conclude

$$\lim_{M \rightarrow \infty} \int_0^M J_0(x) \, dx = 1.$$

References

1. D. Gottlieb and S. Orsag, *Numerical Analysis of Spectral Methods*, SIAM, 1977.
2. M. Pinsky, Fourier inversion for piecewise smooth functions in several variables, *Proceedings of the American Mathematical Society*, 118(1993), 903-910.
3. M. Pinsky, Pointwise Fourier inversion and related eigenfunction expansions, *Communications of Pure and Applied Mathematics*, 47(1994), 653-681.